

(19) World Intellectual Property Organization
International Bureau



FI

(43) International Publication Date
31 July 2003 (31.07.2003)

PCT

(10) International Publication Number
WO 03/062225 A1

(51) International Patent Classification⁷: **C07D 401/12**,
239/48, 413/12, 403/12, 413/14, 409/14, 471/04, A61K
31/506

(74) Agents: **SHUBIN, Harry, B.** et al.; Millen, White, Ze-
lano & Branigan, P.C., Arlington Courthouse Plaza 1, 2200
Clarendon Boulevard, Suite 1400, Arlington, VA 22201
(US).

(21) International Application Number: PCT/US03/01839

(22) International Filing Date: 23 January 2003 (23.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/349,987 23 January 2002 (23.01.2002) US

(71) Applicant (for all designated States except US): **BAYER CORPORATION** [US/US]; 100 Bayer Road, Pittsburgh, PA 15205 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **NAGARATHNAM, Dhanapalan** [IN/US]; 52 Virginia Drive, Bethany, CT 06524 (US). **DUMAS, Jacques** [FR/US]; 98 Farmview Road, Bethany, CT 06524 (US). **HATOUM-MOKDAD, Holia** [US/US]; 43 Joseph Lane, Hamden, CT 06514 (US). **BOYER, Stephen** [US/US]; 233 Colony Street, Fairfield, CT 06430 (US). **WANG, Chunguang** [CN/US]; 27 Timberwood Trail, Hamden, CT 06514 (US). **PLUEMPE, Hans** [DE/DE]; Kirshbaumstrasse 9, D-42115 Wuppertal (DE). **FEURER, Achim** [DE/DE]; Erlbrunnenweg 5, 69259 Wilhelmsfeld (DE). **BENNABI, Samir** [FR/DE]; Alsenstrabe 23 A, 42103 Wuppertal (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIMIDINE DERIVATIVES AS RHO-KINASE INHIBITORS



(1)

(57) Abstract: Disclosed are compounds and derivatives thereof, their synthesis, and their use as Rho-kinase inhibitors. These compounds are useful for inhibiting tumor growth, treating erectile dysfunction, and treating other indications mediated by Rho-kinase, e.g., coronary heart disease.

WO 03/062225 A1

PYRIMIDINE DERIVATIVES AS RHO-KINASE INHIBITORS

Field of the Invention

The present invention relates to compounds and derivatives thereof, their synthesis, and their use as Rho-kinase inhibitors. These compounds of the present invention are useful for inhibiting tumor growth, treating erectile dysfunction, and treating other indications mediated by Rho-kinase, e.g., coronary heart disease.

Background

The pathology of a number of human and animal diseases including hypertension, erectile dysfunction, coronary cerebral circulatory impairments, neurodegenerative disorders and cancer can be linked directly to changes in the actin cytoskeleton. These diseases pose a serious unmet medical need. The actin cytoskeleton is composed of a meshwork of actin filaments and actin-binding proteins found in all eukaryotic cells. In smooth muscle cells the assembly and disassembly of the actin cytoskeleton is the primary motor force responsible for smooth muscle contraction and relaxation. In non-muscle cells, dynamic rearrangements of the actin cytoskeleton are responsible for regulating cell morphology, cell motility, actin stress fiber formation, cell adhesion and specialized cellular functions such as neurite retraction, phagocytosis or cytokinesis (Van Aelst, et al. *Genes Dev* 1997, 11, 2295).

The actin cytoskeleton is controlled by a family of proteins that are a subset of the Ras superfamily of GTPases. This subset currently consists of RhoA through E and RhoG (referred to collectively as Rho), Rac 1 and 2, Cdc42Hs and G25K and TC10 isoforms (Mackay, et al. *J Biol Chem* 1998, 273, 20685). These proteins are GTP (guanine nucleotide triphosphate) binding proteins with intrinsic GTPase activity. They act as molecular switches and cycles between inactive GDP (guanine nucleotide diphosphate) bound and active GTP bound states. Using biochemical and genetic manipulations, it has been possible to assign functions to each family member. Upon activation the Rho proteins controls the formation of actin stress fibers, thick bundles of actin filaments, and the clustering of integrins at focal adhesion complexes. When activated the Rac proteins control the formation of lamellopodia or membrane ruffles on the cell surface and Cdc42 controls filopodia formation. Together this family of proteins plays a critical part in the

control of key cellular functions including cell movement, axonal guidance, cytokinesis, and changes in cell morphology, shape and polarity.

Depending on the cell type and the activating receptor, the Rho proteins can control different biological responses. In smooth muscle cells, Rho proteins are responsible for the calcium sensitization during smooth muscle contraction. In non-smooth muscle cells the Rho GTPases are responsible for the cellular responses to agonist such as lysophosphatidic acid (LPA), thrombin and thromboxane A₂ (Fukata, et al. *Trends Pharmacol Sci* 2001, 22, 32). Agonist response is coupled through heterotrimeric G proteins G_{α12} or G_{α13} (Goetzl, et al. *Cancer Res* 1999, 59, 4732; Buhl, et al. *J Biol Chem* 1995, 270, 24631) though other receptors may be involved. Upon activation Rho GTPases activate a number of downstream effectors including PIP5-kinase, Rhothekin, Rhophilin, PKN and Rho kinase isoforms ROCK-1/ROKβ and ROCK-1/ROKα (Mackay and Hall *J Biol Chem* 1998, 273, 20685; Aspenstrom *Curr Opin Cell Biol* 1999, 11, 95; Amano, et al. *Exp Cell Res* 2000, 261, 44).

Rho kinase was identified as a RhoA interacting protein isolated from bovine brain (Matsui, et al. *Embo J* 1996, 15, 2208). It is a member of the myotonic dystrophy family of protein kinase and contains a serine/threonine kinase domain at the amino terminus, a coiled-coil domain in the central region and a Rho interaction domain at the carboxy terminus (Amano, et al. *Exp Cell Res* 2000, 261, 44). Its kinase activity is enhanced upon binding to GTP-bound RhoA and when introduced into cells, it can reproduce many of the activities of activated RhoA. In smooth muscle cells Rho kinase mediates calcium sensitization and smooth muscle contraction and inhibition of Rho kinase blocks 5-HT and phenylephrine agonist induced muscle contraction. When introduced into non-smooth muscle cells, Rho kinase induces stress fiber formation and is required for the cellular transformation mediated by RhoA (Sahai, et al. *Curr Biol* 1999, 9, 136). Rho kinase regulates a number of downstream proteins through phosphorylation, including myosin light chain (Somlyo, et al. *J Physiol (Lond)* 2000, 522 Pt 2, 177), the myosin light chain phosphatase binding subunit (Fukata, et al. *J Cell Biol* 1998, 141, 409) and LIM-kinase 2 (Sumi, et al. *J Bio Chem* 2001, 276, 670).

Inhibition of Rho kinase activity in animal models has demonstrated a number of benefits of Rho kinase inhibitors for the treatment of human diseases. Several patents

have appeared claiming (+)-trans-4-(1-aminoethyl)-1-(pyridin-4-ylaminocarbonyl)cyclohexane dihydrochloride monohydrate (WO-00078351, WO-00057913) and substituted isoquinolinesulfonyl (EP-00187371) compounds as Rho kinase inhibitors with activity in animal models. These include models of cardiovascular diseases such as hypertension (Uehata, et al. *Nature* 1997, 389, 990), atherosclerosis (Retzer, et al. *FEBS Lett* 2000, 466, 70), restenosis (Eto, et al. *Am J Physiol Heart Circ Physiol* 2000, 278, H1744; Negoro, et al. *Biochem Biophys Res Commun* 1999, 262, 211), cerebral ischemia (Uehata, et al. *Nature* 1997, 389, 990; Seasholtz, et al. *Circ Res* 1999, 84, 1186; Hitomi, et al. *Life Sci* 2000, 67, 1929; Yamamoto, et al. *J Cardiovasc Pharmacol* 2000, 35, 203), cerebral vasospasm (Sato, et al. *Circ Res* 2000, 87, 195; Kim, et al. *Neurosurgery* 2000, 46, 440), penile erectile dysfunction (Chitale, et al. *Nat Med* 2001, 7, 119), central nervous system disorders such as neuronal degeneration and spinal cord injury (Hara, et al. *J Neurosurg* 2000, 93, 94; Toshima, et al. *Stroke* 2000, 31, 2245) and in neoplasias where inhibition of Rho kinase has been shown to inhibit tumor cell growth and metastasis (Itoh, et al. *Nat Med* 1999, 5, 221; Somlyo, et al. *Biochem Biophys Res Commun* 2000, 269, 652), angiogenesis (Uchida, et al. *Biochem Biophys Res Commun* 2000, 269, 633; Gingras, et al. *Biochem J* 2000, 348 Pt 2, 273), arterial thrombotic disorders such as platelet aggregation (Klages, et al. *J Cell Biol* 1999, 144, 745; Retzer, et al. *Cell Signal* 2000, 12, 645) and leukocyte aggregation (Kawaguchi, et al. *Eur J Pharmacol* 2000, 403, 203; Sanchez-Madrid, et al. *Embo J* 1999, 18, 501), asthma (Setoguchi, et al. *Br J Pharmacol* 2001, 132, 111; Nakahara, et al. *Eur J Pharmacol* 2000, 389, 103), regulation of intraocular pressure (Honjo, et al. *Invest Ophthalmol Vis Sci* 2001, 42, 137) and bone resorption (Chellaiah, et al. *J Biol Chem* 2000, 275, 11993; Zhang, et al. *J Cell Sci* 1995, 108, 2285).

The inhibition of Rho kinase activity in patients has benefits for controlling cerebral vasospasms and ischemia following subarachnoid hemorrhage (*Pharma Japan* 1995, 1470, 16).


Summary of the Invention

The compounds and their derivatives presented in this invention are useful as Rho Kinase inhibitors and thus have utilities in the treatment of cardiovascular disease, e.g.,

hypertension, atherosclerosis, restenosis, cerebral ischemia, and cerebral vasospasm, as well as neuronal degeneration, spinal cord injury, cancers of the breast, colon, prostate, ovaries, brain and lung and their metastases, thrombotic disorders, asthma, glaucoma and osteoporosis.

In addition, the compounds of the invention are useful to treat erectile dysfunction, i.e., erectile dysfunction mediated by Rho-kinase. Erectile dysfunction can be defined as an inability to obtain or sustain an erection adequate for intercourse, WO 94/28902, U.S.P. 6,103,765 and U.S.P. 6,124,461.

The invention involves compounds of formula I




wherein R_1 and R_2 are each independently

H, halo, alkyl, optionally substituted by halo up to the perhalo level, cycloalkyl, alkenyl, alkynyl, NO_2 , NH_2 , NR_6R_7 , or furyl, thienyl, pyridyl, trifluoromethyl or phenyl each optionally substituted with NH_2 , NO_2 , trifluoromethyl or alkoxy;

R_1 and R_2 may be taken together to form a ring of from 5 to 7 members optionally interrupted by N and optionally substituted on N by benzyl;

R_3 is NH_2 or $-\text{NH}-$ phenyl optionally substituted with halo, C_1-C_4 alkyl, trifluoromethyl, nitro or amino;

R_4 is



X is a linker substituted at the 3 or 4 position of the ring and is selected from

O, S, $-\text{S}-\text{CH}_2-$, $-(\text{CH}_2)_m-$, or $-(\text{C}=\text{O})-$;

A is phenyl optionally substituted with alkylthio or OH,
pyridyl,
quinolyl or

isoquinolyl;

each R_5 independently is halo, alkyl optionally substituted by halo up to the perhalo level, cycloalkyl, alkoxy, alkenyl, alkynyl, NO_2 , NH_2 , or trifluoromethyl;



n is 0, 1, 2, 3 or 4;





m is 1 or 2; and




R_6 and R_7 are each independently H, alkyl, cycloalkyl, or phenyl optionally substituted with halo, CF_3 , alkyl, nitro or amino; or





R_6 and R_7 may form, together with the N atom to which they are attached, a heterocyclic ring optionally substituted with alkyl, optionally interrupted by O, or optionally fused to phenyl;





with the proviso that formula I does not include the following compounds:


























































Preferred compounds of formula I include those wherein

R_1 and R_2 are each independently H, halo, C_{1-12} alkyl optionally substituted by halo up to the perhalo level, C_{2-12} , alkenyl, C_{2-12} -alkynyl, NO_2 , NH_2 , NR_6R_7 , alkyl, furyl, thienyl, pyridyl, phenyl optionally substituted with NO_2 , trifluoromethyl or NR_6R_7 ;

R_1 and R_2 may be taken together to form a ring of from 5 to 7 members optionally interrupted by N and (optionally) substituted on N by benzyl;

R_3 is NH_2 - or $-NH-$ phenyl optionally substituted with halo, C_1-C_4 alkyl, trifluoromethyl, nitro or amino;

R_4 is



X is a linker substituted at the 3 or 4 position of the ring and is selected from

O, S, $-S-CH_2-$, $-CH_2-S-$, $-(CH_2)_m-$, or $-(C=O)-$;

A is phenyl optionally substituted by C_{1-4} alkylthio or OH,

pyridyl,

quinolyl or

isoquinolyl;

each R_5 independently is halo, C_1-C_{12} alkyl, optionally substituted by halo, up to the perhalo level, C_{2-12} -alkenyl, C_{2-12} -alkynyl, NO_2 , NH_2 or trifluoromethyl;


n is 0, 1, 2, 3 or 4;

m is 1 or 2; and

R₆ and R₇ are each independently H, C₁-C₆ alkyl, or phenyl optionally substituted with halo, CF₃, C₁-C₄ alkyl, nitro or amino, with the exception of the compounds previously cited.

Particularly preferred compounds include those wherein

- (i) X is not S or A is not pyridyl, or both;
- (ii) X is not S or -S-CH₂-;
- (iii) n is 1-4
- (iv) n is 1-4 and each R₅ is independently is halo, C₃-C₁₂ alkyl optionally substituted by halo up to the perhalo level, C₂-₁₂-alkenyl, C₂-₁₂-alkynyl, NO₂, NH₂ or trifluoromethyl; or
- (v) the compound of formula I is not



The terms identified above have the following meaning throughout:

“Alkyl” means straight or branched chain alkyl groups having from one to about twelve carbon atoms. Such groups include methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, *neo*-pentyl, 2-pentyl, *n*-hexyl, 2-hexyl, 3-hexyl, dodecyl, 2,3-dimethylbutyl and the like.

“Cycloalkyl” means saturated monocyclic alkyl groups of from 3 to about 8 carbon atoms and includes such groups as cyclopropyl, cyclopentyl, cyclohexyl and the like.

“Alkenyl” means straight or branched chain alkenyl groups having from two to about twelve carbon atoms. Such groups include vinyl, allyl, isopropenyl, 3-butenyl and the like.

"Alkynyl" means straight or branched chain alkynyl groups having from two to about twelve carbon atoms. Such groups include ethynyl, propargyl, 3-pentynyl, 3-heptynyl, 1-methyl-2-butyne and the like.

"Alkoxy" means straight or branched chain alkoxy groups having from about one to eight carbon atoms and includes such groups as methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, isobutoxy, *sec*-butoxy, *tert*-butoxy and the like.

"Halo" means fluoro, chloro, bromo, or iodo. Preferred are fluoro, chloro and bromo, more preferably fluoro and chloro.

When an alkyl substituent is described as being substituted by oxo, it means substitution by a doubly bonded oxygen atom, which forms together with the carbon to which it is attached, a carbonyl group $-(C=O)-$.

The term "optionally substituted" means that the moiety so modified may be unsubstituted or substituted with the identified substituent(s).

When any moiety is described as being substituted, it can have one or more of the indicated substituents that can be located at any available position on the moiety. When there are two or more substituents on any moiety, each substituent is defined independently of any other substituent and can, accordingly, be the same or different.

The compounds of the Formula I can be made according to routine, conventional chemical methods, and/or as disclosed below, from starting materials which are either commercially available or producible according to routine, conventional chemical methods. General methods for the preparation of the compounds are given below, and the preparation of representative compounds is specifically illustrated in the Examples.


General Methods of Preparation

Compounds of Formula I may be prepared using one of the general methods summarized below in Reaction Schemes 1-3, from either commercially available or readily prepared starting materials. Preparation of starting material is described in Reaction Schemes 4-6.

In the first methods, illustrated in Reaction Scheme 1, the reaction of a chloropyrimidine of Formula II with a substituted aromatic compound of Formula III,


where R₁–R₄ are as defined above, may be accomplished under either basic or acidic conditions (Reaction Scheme 1).

Reaction Scheme 1




Alternatively, when R₁ is an aryl group, a two-step scheme may be employed. In the first step, the Formula III compound is allowed to react with the dichloropyrimidine of Formula IIa under acidic conditions as in Reaction Scheme 1. In the second step, the R₁ group is introduced by palladium-catalyzed coupling of a boronic acid of Formula R₁B(OH)₂ (e.g., Suzuki coupling) to provide compounds of Formula I in which R₁ is aryl (Reaction Scheme 2).

Reaction Scheme 2




The compounds of Formula I, where R₁ is NR₆R₇, may be prepared conveniently by reaction the chloropyrimidine of Formula Ia with a variety of amines of Formula R₆R₇NH, usually in a higher boiling polar solvent such as *n*-butanol, and at an elevated temperature, e.g., 90-120 °C as shown in Reaction Scheme 3.

Reaction Scheme 3




General Method of Preparation of Intermediates

Intermediate chloropyrimidines of Formula II and aryl amines of Formula III are either commercially available or may be prepared by methods well known in the art. For example, a pyrimidone can be converted to the corresponding chloropyrimidine II by reaction with POCl_3 , and reduction of nitro aromatic compounds under standard conditions (e.g., Fe, HOAc) provides aryl amines of Formula III. Further illustrative methods are depicted in Reaction Schemes 4–6 below. For example, in Reaction Scheme 4, Formula III compounds in which R_4 is




, A is 2 or 4 pyridyl, and X is O or S, are prepared by a nucleophilic aromatic substitution reaction of a haloarene (A-halo) with a phenol or thiophenol of Formula IV, carried out in a polar solvent such as DMF and assisted by a base such as potassium carbonate (Reaction Scheme 4).

Reaction Scheme 4



Intermediates of Formula IIIb may be prepared as shown in Reaction Scheme 5 in two steps from a nitroaromatic compound of Formula V, where Z represents either CH or N. The nitro aromatic compound is allowed to react in basic media at elevated temperatures, and the resulting nitro aromatic compound is reduced by standard means, e.g., iron and acetic acid.


Reaction Scheme 5



a) base, polar solvent e.g., DMF, heat
 b) reduction, e.g., Fe, HOAc, H₂O, rt

A wide variety of chloropyrimidines of Formula II where R₃ is NH₂ and R₁ is other than NR₆R₇, may be prepared by the route shown in Reaction Scheme 6 below. Reaction of a carboxylic acid of Formula VI, where R₁ is other than NR₆R₇, with a substituted Meldrum's acid of Formula VII (prepared, for example by alkylation of Meldrum's acid with R₂-halo and base), gives a ketoester of Formula VIII. The pyrimidone compound of Formula IX is formed by reaction of compound VIII with guanidine in the presence of an acid in protic solvent such as ethanol. The pyrimidone of Formula IX is then converted to the chloropyrimidine IIb, (II, where R₃ is NH₂) under typical conditions, e.g., POCl₃ and heat.

Reaction Scheme 6



R'' = lower alkyl, benzyl

a) DMAP/DCC/CH₂Cl₂, 0 °C - rt; b) pTsOH.H₂O/R''OH, reflux; c) Guanidine carbonate, HCl, EtOH, reflux; d) POCl₃, 100 °C

Without further elaboration, it is believed that one skilled in the art can, using the preceding descriptions, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius; and, unless otherwise indicated, all parts and percentages are by weight.

The entire disclosure of all applications, patents and publications, cited above or below, and U.S. Provisional Application Serial No. 60/349,987, filed January 23, 2002, are hereby incorporated by reference.

Abbreviations and Acronyms

When the following abbreviations are used herein, they have the following meaning:

Ac ₂ O	acetic anhydride
anhy.	anhydrous
<i>n</i> -BuOH	<i>n</i> -butanol
<i>t</i> -BuOH	<i>t</i> -butanol
CD ₃ OD	methanol- <i>d</i> ₄
Celite®	diatomaceous earth filter agent, ® Celite Corp.
CH ₂ Cl ₂	methylene chloride
CI-MS	chemical ionization mass spectroscopy
conc	concentrated
dec	decomposition
DME	dimethoxyethane
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
ELSD	evaporative light scattering detector
EtOAc	ethyl acetate
EtOH	ethanol (100%)
Et ₂ O	diethyl ether
Et ₃ N	triethylamine
HPLC ES-MS	high performance liquid chromatography-electrospray mass spectroscopy
LC	liquid chromatography
MPLC	medium pressure liquid chromatography
MS	mass spectrometry

NMM	4-methylmorpholine
Ph ₃ P	triphenylphosphine
Pd(dppf)Cl ₂	[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II)
Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium(0)
Pd(OAc) ₂	palladium acetate
P(O)Cl ₃	phosphorous oxychloride
RT	retention time (HPLC0)
rt	room temperature
THF	tetrahydrofuran
TFA	trifluoroacetic acid
TLC	thin layer chromatography

Experimental Examples


All reactions were performed in flame-dried or oven-dried glassware under a positive pressure of dry argon, and were stirred magnetically unless otherwise indicated. Sensitive liquids and solutions were transferred via syringe or cannula, and introduced into reaction vessels through rubber septa. Commercial grade reagents and solvents were used without further purification. Thin layer chromatography (TLC) was performed on Analtech UNIPLATE™ pre-coated glass-backed silica gel 60 A F-254 250 μm plates. Column chromatography (flash chromatography) was performed on a Biotage system using 32-63 micron, 60 A, silica gel pre-packed cartridges. Proton (¹H) nuclear magnetic resonance (NMR) spectra were measured with a Varian (300 MHz) spectrometer with residual protonated solvent (CHCl₃ δ 7.26; MeOH δ 3.30; DMSO δ 2.49) as standard. Low-resolution mass spectra (MS) were either obtained as electron impact (EI) mass spectra or as fast atom bombardment (FAB) mass spectra.

The IUPAC name was obtained using the ACD/ILab Web service.


A. Preparation of chloropyrimidine intermediates

Intermediate A1

Preparation of 2-amino-4-chloro-5,6-dimethyl-pyrimidine




Step 1. Preparation of 2-amino-5,6-dimethyl-4-pyrimidinone



To a solution of ethyl 2-acetoacetate (6.0 g, 41.6 mmol) and guanidine carbonate (5.6 g, 31.2 mmol) in EtOH (32 mL) was added 12 N HCl (350 μ L). The mixture was refluxed for 16 h. After the reaction was cooled to room temperature, the solid was collected by filtration and washed with EtOH. A solution of the solid in 1 N NaOH was refluxed for 3 h. After the reaction was cooled to room temperature, the aqueous mixture was adjusted to pH = 5 with concentrated acetic acid. The resulting precipitate was collected by filtration, washed with water and then with hexanes, and dried under vacuum. Desired compound (6.34 g, 45.6 mmol; 100% yield); ^1H NMR (D_2O ; NaOD) δ 1.47 (s, 3H), 1.29-1.30 (m, 2H), 1.22 (s, 3H); ES MS $[\text{M}+\text{H}]^+ = 140$.

Step 2. Preparation of 2-amino-4-chloro-5,6-dimethyl-pyrimidine




The product of the previous step (2.0 g, 14.4 mmol) and phosphorus oxychloride (6 mL, 57.5 mmol), was refluxed for 4 h. The reaction was cooled to rt and poured over ice. The mixture was separated and the aqueous layer was extracted with chloroform (3 x 75 mL). The aqueous mixture was adjusted to pH = 9 with concentrated ammonium hydroxide. The resulting solid product was collected by filtration, washed with water, and


dried under vacuum. Desired compound (963 mg, 6.1 mmol; 43% yield); mp = 212 - 220°C; ES MS $[M+H]^+ = 158$; TLC (CH_2Cl_2 -MeOH, 90:10); $R_f = 0.72$.

Intermediate A2

Preparation of 2-amino-4-chloro-6-(4-pyridyl)pyrimidine




Step 1. Preparation of 2-amino-4-hydroxy-6-(4-pyridyl)pyrimidine



A solution of guanidine carbonate (7.1 g, 39 mmol, 1.5 eq), ethyl isonicotinoyl acetate (10 g, 51.76 mmol), and hydrochloric acid (0.75 mL, 9.0 mmol) in absolute ethanol (80 mL) was refluxed under argon overnight. The precipitate formed was filtered, washed with ethanol and dried. The solid was then dissolved in 1 N NaOH (100 mL) and refluxed for 2 h. The reaction mixture was then cooled to room temperature, acidified with glacial acetic acid, and the solid formed was filtered and dried to afford the desired product as a white solid (5.45 g, 56%). $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ 6.24 (s, 1H), 6.79 (bs, 2H), 7.85 (d, $J=5.1$ Hz, 2H), 8.62 (d, $J=5.3$ Hz, 2H), 11.22 (bs, 1H).

Step 2. Preparation of 2-amino-4-chloro-6-(4-pyridyl)pyrimidine :




A solution of 2-amino-4-hydroxy-6-(4-pyridyl)pyrimidine (5.45 g, 29 mmol) in POCl_3 (12 mL) was refluxed under argon for 5 h. The reaction mixture was cooled to room temperature, poured over ice, and allowed to stir at room temperature for 2 h to ensure the quenching of POCl_3 . At this time, the mixture was made basic upon addition of 1 N NaOH and the brown solid was filtered to afford 4.52 g of crude product, which


was used without further purification (NMR analysis showed 1:1 product / starting material). The filtrate formed more solid upon standing at room temperature (1 g, NMR analysis showed 2:1 product / starting material). ^1H -NMR (DMSO- d_6) δ 7.34 (bs, 2H), 7.38 (s, 1H), 7.99 (d, $J=4.2$ Hz, 2H), 8.72 (d, $J=4.6$ Hz, 2H).

Intermediate A3


Preparation of 2-amino-4-chloro-6-(2-thienyl)pyrimidine




Step 1. Preparation of ethyl-2-(thiophene-2-yl)acetate.





A solution of thiophene-2-carboxylic acid (8.9 g, 68.5 mmol), 2,2-dimethyl-1,3-dioxane-4,6-dione (12.0 g, 81.6 mmol), and 4-dimethylaminopyridine (17.0 g, 138 mmol) in dry CH_2Cl_2 (100 mL) was cooled to 0 °C and treated with a solution of 1,3-dicyclohexylcarbodiimide (75 mL, 1.0 M in CH_2Cl_2 , 75 mmol). The reaction was allowed to stir at room temperature for 2 h and the dicyclohexylurea was then filtered and washed with CH_2Cl_2 . The filtrate was concentrated at reduced pressure and the residue was dissolved in absolute ethanol (400 mL). The solution was then treated with a solution of *p*-toluenesulfonic acid monohydrate (32 g, 168 mmol) in absolute ethanol (100 mL) and refluxed under argon for 1 h. At this time, the ethanol was removed at reduced pressure and the residue was dissolved in EtOAc and washed sequentially with H_2O (300 mL), saturated NaHCO_3 (200 mL), 1 N HCl (200 mL), saturated NaCl, and dried (MgSO_4). The solvent was removed at reduced pressure and the residue was filtered through a pad of silica with 10% EtOAc/90% hexanes to afford the desired product as an oil (13 g, 96%). TLC (20% EtOAc/80% hexane) R_f 0.21; ^1H -NMR (DMSO- d_6) δ 1.17 (t, $J=7.01$, 3H), 4.06-4.14 (m, 4H), 7.25 (t, $J=5.1$ Hz, 1H), 7.98 (d, $J=3.8$ Hz, 1H), 8.06 (d, $J=4.9$ Hz, 1H).

Step 2. Preparation of 2-amino-4-hydroxy-6-(2-thienyl)pyrimidine.


The procedure was similar to that used for Intermediate A2, step 1, using ethyl-2-(thienyl-2-yl)acetate as starting material. (43% yield). TLC (6% MeOH/94% CH₂Cl₂) R_f 0.23; MS ES 194 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 6.06 (s, 1H), 6.70 (bs, 2H), 7.11 (t, *J*=4.9 Hz, 1H), 7.64 (d, *J*=4.9 Hz, 1H), 7.70 (d, *J*=3.6 Hz, 1H), 10.95 (bs, 1H).

Step 3. Preparation of 2-amino-4-chloro-6-(2-thienyl)pyrimidine.



The procedure was similar to that of Intermediate A2, step 2, using 2-amino-4-hydroxy-6-(2-thiophene)pyrimidine as starting material. It afforded 33% yield after purification on silica with 15% EtOAc/85% hexanes. TLC (20% EtOAc/80% hexanes) R_f 0.29; ¹H-NMR (DMSO-*d*₆) δ 7.16-7.23 (m, 4H), 7.77 (dd, *J*=0.8, 5.0 Hz, 1H), 7.98 (dd, *J*=1.0, 3.8 Hz, 1H).

Intermediate A4Preparation of 2-amino-4-chloro-6-(2-furyl)pyrimidineStep 1. Preparation of 2-amino-4-hydroxy-6-(2-furyl)pyrimidine.

The general procedure for the preparation of Intermediate A2, (step 1) was used; (37% yield). MS (ES) 178 [M+H]⁺.


Step 2. Preparation of 2-amino-4-chloro-6-(2-furyl)pyrimidine.

A solution of 2-amino-4-hydroxy-6-(2-furyl)pyrimidine (1.40 g, 7.9 mmol) in POCl₃ (4 mL) was refluxed under argon for 2 h. The POCl₃ was distilled; the residue was diluted with EtOAc and poured over iced saturated NaHCO₃. The layers were separated and the aqueous was extracted with EtOAc (100 mL). The combined extracts was washed with saturated NaCl, dried (MgSO₄), and the solvent removed at reduced pressure to afford 0.5 g of crude product, which was used without further purification. TLC (20% EtOAc/80% hexane) R_f 0.26; ¹H-NMR (DMSO-*d*₆) δ 6.68 (dd, *J*=1.7, 3.4 Hz, 1H), 6.94 (s, 1H), 7.25 (dd, *J*=1, 3.7 Hz, 1H), 7.91 (dd, *J*=0.8, 1.9 Hz, 1H).

Intermediate A5Preparation of 6-benzyl-4-chloro-5,6,7,8-tetrahydropyrido[4,3-*d*]pyrimidin-2-amineStep 1. Preparation of 2-amino-7-benzyl-5,6,7,8-tetrahydropyrido[3,4-*d*]pyrimidin-4(3H)-one.

To EtOH (16 mL) cooled to 0 °C (ice/H₂O bath) was added Na spheres (204 mg, 8.9 mmol). The mixture was stirred until all Na dissolved. Methyl 1-benzyl-4-oxo-3-piperidine-carboxylate hydrochloride (3.0 g, 10.1 mmol) and guanidine carbonate (1.4 g, 7.6 mmol) were added. The mixture was refluxed for 16 h. After the reaction was cooled to room temperature, the solid was collected by filtration, washed with EtOH, and dried under vacuum. Desired compound (2.58 g, 10.0 mmol; 99+% yield); mp = 202 - 212 ° (dec.); ES MS [M+H]⁺ = 257; TLC (CH₂Cl₂-MeOH, 90:10); R_f = 0.20.


Step 2. Preparation of 6-benzyl-4-chloro-5,6,7,8-tetrahydropyrido[4,3-*a*]pyrimidin-2-amine.



A solution of the product from step 1 (3.5 g, 13.7 mmol) in POCl_3 (52 mL) was refluxed under argon for 5 h. The reaction mixture was cooled to room temperature, poured over ice, and allowed to stir at room temperature for 2 h to ensure the quenching of POCl_3 . At this time, the mixture was made basic upon addition of ammonium hydroxide and was extracted with CH_2Cl_2 (3 x 200 mL). The combined organics were washed with 1N NaOH followed by brine, dried (MgSO_4), and concentrated under reduced pressure. The residue was taken up in benzene and was made acidic upon the addition of 1N HCl in diethyl ether. The brown solid was filtered to afford 0.35 g of crude product, which was used without further purification. ES MS $[\text{M}+\text{H}]^+ = 275$.

Intermediate A6


Preparation of 2-amino-6-(trifluoromethyl)-4-pyrimidinyl 4-methylbenzenesulfonate





To a solution of 2-amino-6-(trifluoromethyl)-4(3*H*)-pyrimidinone (250 mg, 1.4 mmol), triethylamine (196 μL , 1.4 mmol), *N,N*-dimethylaminopyridine (17 mg, 0.14 mmol), in CH_2Cl_2 (13 mL) cooled to 0 °C was added *p*-toluenesulfonyl chloride (534 mg, 2.8 mmol). The mixture was stirred at room temperature for 16 h. The mixture was diluted with CH_2Cl_2 , washed with H_2O (2x 20 mL) followed by brine, dried (Na_2SO_4), evaporated, and dried under vacuum. Desired compound (466 mg, 1.4 mmol; 99+% yield; ES MS $[\text{M}+\text{H}]^+ = 140$).

Using the above methods for the preparation of A1-A6 and substituting the appropriate starting materials, the following pyrimidine intermediates were also prepared.


Table 1
Chloropyrimidine Intermediates A



<u>Intermediate No.</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	<u>Physical Data</u>
A7	Me	H	NH ₂	Aldrich
A8	Et	H	NH ₂	Aldrich or Lancaster
A9	H	H	NH ₂	Aldrich
A10	<i>t</i> -Bu	H	NH ₂	mp = 109-113 °C; ES MS [M+H] ⁺ =186; TLC (90:10 CH ₂ Cl ₂ /MeOH); R _f = 0.37.
A11	Me	Cl	NH ₂	Aldrich or Lancaster?
A12	-(CH) ₄ -		NH ₂	¹ H NMR (DMSO- <i>d</i> ₆) δ 6.60 (s, 2H), 2.55-2.29 (m, 4H), 1.68-1.56 (m, 4H)
A13	-(CH) ₅ -		NH ₂	¹ H NMR (DMSO- <i>d</i> ₆) δ 6.65 (s, 2H), 2.72-2.58 (m, 4H), 1.74-1.60 (m, 2H), 1.55-1.34 (m, 4H)
A14	-(CH) ₃ -		NH ₂	¹ H NMR (DMSO- <i>d</i> ₆) δ 6.73 (s, 2H), 2.72-2.57 (m, 4H), 1.89 (sept, <i>J</i> = 7.0, 2H)
A15	<i>i</i> -Pr	H	NH ₂	mp = 104-112 °C; ¹ H NMR (D ₂ O) δ 6.11 (s, 1H), 2.23-2.11 (m, 1H), 0.46 (d, <i>J</i> = 6.2 Hz, 6H); ES MS [M+H] ⁺ = 172
A16	CH ₃	H	Ph-NH-	
A17	Ph	H	NH ₂	
A18	3-pyridyl	H	NH ₂	
A19	2-pyridyl	H	NH ₂	
A20	3-NO ₂ -Ph	H	NH ₂	
A21	Cl	H	NH ₂	Aldrich

B. Preparation of arylamine intermediates**Intermediate B1****Preparation of 1-(4-pyridinyl)-1H-indol-5-amine****Step 1. Preparation of 5-nitro-1-(4-pyridinyl)-1H-indole**

To a solution of 5-nitroindole (7.0 g, 43.2 mmol) and 4-chloropyridine hydrochloride (7.8 g, 51.8 mmol) in DMF (43 mL) was added potassium *tert*-butoxide (12.1 g, 108.0 mmol), portionwise. The reaction was heated at 100 °C for 48 h. The mixture was allowed to cool to room temperature and poured into water (400 mL). The resulting solid was removed by filtration and dried under vacuum. Desired compound (6.04 g, 25.3 mmol; 58% yield); ¹H NMR (DMSO-*d*₆) δ 8.76 (dd, *J* = 1.7, 4.5, 2H), 8.68 (d, *J* = 2.2, 1H), 8.06-8.13 (m, 2H), 7.92 (d, *J* = 9.2, 1H), 7.75 (dd, *J* = 1.5, 4.6, 2H), 7.07 (dd, *J* = 0.9, 3.5, 1H); ES MS [M+H]⁺ = 240.


Step 2. Preparation of 1-(4-pyridinyl)-1H-indol-5-amine

A mixture of the product from step 1 (8.27 g, 34.6 mmol) and 10% palladium-on-charcoal catalyst (827 mg) in ethyl acetate (166 mL) and EtOH (9 mL) was stirred under hydrogen at atmospheric pressure for 48 h. Further catalyst (414 mg) was added and the reaction was stirred for 24 h. Again, further catalyst (414 mg) was added and the reaction was stirred an additional 24 h. The catalyst was removed by filtration through


diatomaceous earth and the solvent removed from the filtrate by evaporation. The residue was triturated with ether, collected by filtration, and dried under vacuum. Desired compound (4.67 g, 22.3 mmol; 65% yield); mp = 149 - 154 °C; ES MS $[M+H]^+ = 210$; TLC (CH_2Cl_2 -MeOH, 95:5); $R_f = 0.29$.

Intermediate B2

Preparation of 4-[(4-aminophenyl)sulfanyl]phenol




Step 1. Preparation of 4-[(4-nitrophenyl)sulfanyl]phenol.



To a solution of nitrobenzenesulfonyl chloride (4g, 21 mmol) in ether (25 mL) was added phenol (1.97 g, 20 mmol) as a solution in ether (25 mL). After being stirred for 15 h at rt, the mixture was concentrated to afford a crude solid which was recrystallized from acetic acid. Desired compound (4.0 g, 16.2 mmol, 76% yield). TLC (Hexanes/EtOAc, 70:30); $R_f = 0.54$.


Step 2. Preparation of 4-[(4-aminophenyl)sulfanyl]phenol.




To a solution of the product of step 1 (4g, 16.2 mmol) in EtOH (500 mL) was added $SnCl_2 \cdot 2H_2O$ (18.3 g, 81 mmol). The solution was warmed to reflux. After being stirred for 3 h, the mixture was allowed to cool to rt, and the volatiles were removed by rotary evaporation. The resultant slurry was suspended in EtOAc, and solid $NaHCO_3$ was added. Subsequently, the mixture was filtered, and the filtered solid was washed thoroughly with EtOAc. The organic filtrate was washed with water, and the aqueous washes were extracted with EtOAc. The combined organic extracts were washed with brine, dried ($MgSO_4$), filtered, and concentrated to afford an orange solid, which was used without additional purification. Desired compound (3.0 g, 13.8 mmol, 86 % yield). TLC (Hexanes/EtOAc, 70:30); $R_f = 0.34$.

Intermediate B3

Preparation of (3-aminophenyl)[4-(methylsulfanyl)phenyl]methanone




Step 1. Preparation of [4-(methylsulfanyl)phenyl](3-nitrophenyl)methanone





3-nitrobenzoylchloride (5.0 g, 26.94 mmol) was added to a solution of thioanisole (3.16 mL, 26.94 mmol) and 1,2-dichloroethane (95 mL). The resulting reaction mixture was cooled to 0 °C (ice/H₂O bath) and 0.5 equivalents of aluminum trichloride (1.8 g, 13.47 mmol) was added. The reaction was allowed to stir for 15 min at this temperature and the cold bath was removed followed by addition of the remaining equivalents of AlCl₃ (2.51 g, 18.87). The reaction solution turned a dark greenish/yellow and was allowed to stir at room temp. for 18h, after which time the reaction was quenched slowly with H₂O (50 mL). The mixture was diluted with CH₂Cl₂ (50 mL) and washed with H₂O (3 x 50 mL), and the combined organic phases were washed with satd NaHCO₃ (50 mL), dried (MgSO₄) and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography (EtOAc/hexane, 1/4) to afford 3.3 g (44%) of 4-(methylsulfanyl)phenyl[(3-nitrophenyl)methanone as a solid. EI-LRMS *m/z* 274 (M⁺); TLC R_f 0.68 (EtOAc/Hex, 2/3).


Step 2. Preparation of (3-aminophenyl)[4-(methylsulfanyl)phenyl]methanone




Prepared analogously to Intermediate B2, step 2. The crude product was purified by flash column chromatography, eluting with 70:30 Hexanes/EtOAc. TLC: (Hexanes/EtOAc, 70:30); R_f 0.15.

Intermediate B4**Preparation of 4-(4-aminophenoxy)phenol****Step 1. Preparation of 4-(4-nitrophenoxy)phenol**

A mixture of *p*-nitrofluorobenzene (25 g, 0.177 mol), dihydroquinone (19.5 g, 0.177 mol), and sodium hydroxide (7.08 g, 0.177 mol) in EtOH/H₂O (1:1 v/v, 176 mL) was heated at reflux for 20 h, and subsequently allowed to cool to room temperature. The mixture was filtered, the filtrate was made acidic with dilute aqueous HCl, and the resultant precipitate filtered to afford the crude product as a yellow solid. The desired product was recrystallized from EtOH. (15 g, 0.064 mol, 37 % yield). . TLC (Hexanes/EtOAc, 70:30); *R*_f = 0.44.

Step 2. Preparation of 4-(4-aminophenoxy)phenol

To a solution of the product of step 1 in EtOH (100 mL) was added 10% palladium on carbon (200 mg). After being stirred under an atmosphere of hydrogen overnight, the mixture was filtered through Celite[®]. The volatiles were removed from the filtrate to provide the crude product which was purified by flash column chromatography eluting with Hexanes/EtOAc (85:15, followed by 75:25). Desired product (1.5 g, 7.45 mmol, 86 %). TLC (Hexanes/EtOAc, 70:30); *R*_f = 0.41.


Intermediate B5**Preparation of 4-(4-pyridinylthio)aniline**

To a solution of 4-aminothiophenol (20.2 g, 156.5 mmol) in anhydrous DMF (200 mL) was added 4-chloropyridine hydrochloride (24.4 g, 161.0 mmol) followed by


potassium carbonate (44 g, 318.4 mmol). The reaction mixture was heated at 80 °C overnight, then diluted with ethyl acetate (400 mL) and water (400 mL). The aqueous layer was back-extracted with ethyl acetate (2 x 200 mL). The combined organic layers were washed with a saturated aqueous NaCl solution (200 mL), dried over anhydrous MgSO_4 , and concentrated under reduced pressure. The residue was filtered through a pad of silica with ethyl acetate and the resulting material was triturated with an ethyl ether / hexane solution to afford the desired product (24.7 g, 78%). TLC (50% ethyl acetate / 50% hexane) R_f 0.25; $^1\text{H-NMR}$ (DMSO-d_6) δ 5.67 (bs, 2H), 6.65 (d, $J=8.4$ Hz, 2H), 6.88 (d, $J=6.2$ Hz, 2H), 7.19 (d, $J=8.4$ Hz, 2H), 8.27 (d, $J=6.2$ Hz, 2H), $\text{MS}[\text{M}+\text{H}]^+ = 203$.

Intermediate B6

Preparation of 4-[2-(4-pyridinyl)ethyl]aniline




Step1. Preparation of 4-[(E)-2-(4-nitrophenyl)ethenyl]pyridine



To an oven dried 500 mL 3-necked flask was added (4-nitrobenzyl)triphenylphosphonium bromide (15 g, 30.42 mmol) followed by the addition of THF (100 mL). The solution was cooled to 0 °C in an ice bath. Potassium *t*-butoxide (3.9 g, 33.02 mmol) was then added in one portion resulting in an orange suspension. The suspension was maintained at 0 °C while a solution of 4-pyridine-2-carboxaldehyde (2.7 g, 24.70 mmol) in THF (20 mL) was added in 10 minutes. The ice bath was removed and the reaction was stirred at room temperature for 2 h. At this time, the reaction was quenched with saturated ammonium chloride solution (50 mL) and stirred for 15 minutes. The mixture was then extracted with ethyl acetate (2 x 100 mL), the combined extracts were washed with saturated aqueous NaCl solution (100 mL) and dried (MgSO_4). The solvent was removed at reduced pressure and the residue was chromatographed on silica with 0-50% ethyl acetate in hexanes to afford the desired product (1.8 g, 32%). TLC (50% ethyl acetate / 50% hexane) R_f 0.28; $^1\text{H-NMR}$ (DMSO-

d_6) δ 6.84 (d, $J=12.4\text{Hz}$, 1H), 6.96 (d, $J=12.4\text{Hz}$, 1H), 7.14 (d, $J=6.2\text{Hz}$, 2H), 7.45 (d, $J=8.7\text{Hz}$, 2H), 8.15 (d, $J=8.7\text{Hz}$, 2H), 8.47 (d, $J=6.2\text{Hz}$, 2H).


Step 2. Preparation of 4-[2-(4-pyridinyl)ethyl]aniline




To a dry 50 mL flask flushed with argon was added 10% Pd on carbon (285 mg) followed by the addition of ethanol (12 mL) and the product from step 1 (1.8 g, 8.0 mmol). At this time, the argon line was replaced with a hydrogen balloon and the reaction was stirred overnight. The mixture was filtered through a pad of Celite[®] and the filtrate was concentrated at reduced pressure. The solid residue was triturated with ethyl ether/hexanes to afford the desired product (1.2 g, 67%). TLC (4% acetone / 96% methylene chloride) R_f 0.09; $^1\text{H-NMR}$ (DMSO- d_6) δ 2.67-2.83 (m, 4H), 4.83 (bs, 2H), 6.45 (d, $J=8.2\text{Hz}$, 2H), 6.84 (d, $J=8.2\text{Hz}$, 2H), 7.20 (d, $J=6\text{Hz}$, 2H), 8.41 (d, $J=6\text{Hz}$, 2H).

Intermediate B7

Preparation of 3-fluoro-4-(4-pyridinylsulfanyl)aniline




Step 1. Preparation of 4-[(2-fluoro-4-nitrophenyl)sulfanyl]pyridine.



A solution of 4-mercaptopyridine (4.2 g, 35.6 mmol), 3,4-difluoronitrobenzene (5.7 g, 35.7 mmol), and potassium carbonate (12.4 g, 89.7 mmol) in anhydrous DMF (40 mL) was stirred at 40 °C and under argon for 3 h. TLC showed complete reaction. The mixture was diluted with ethyl acetate (100 mL) and water (100 mL) and the aqueous layer was back-extracted with ethylacetate (2 x 100 mL). The organic layers were washed with a saturated NaCl solution (100 mL), dried (MgSO₄), and concentrated under reduced pressure. The crude product was purified by column chromatography with 50% ethyl acetate / 50% hexanes. It afforded the desired product as a yellow solid (6.3 g,

71%). TLC (50% EtOAc/50% hexane) R_f 0.53; $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ 7.27 (dd, $J=0.76$, 4.2 Hz, 2H), 7.78 (dt, $J=0.76$, 7.2 Hz, 1H), 8.11-8.15 (m, 1H), 8.28-8.33 (m, 1H), 8.5 (dd, $J=1.4$, 4.6 Hz, 2H), MS $[\text{M}+\text{H}]^+ = 251$.


Step2. Preparation of 3-fluoro-4-(4-pyridinylsulfanyl)aniline.



A slurry of 3-fluoro-4-pyridinylthio)nitrobenzene (6.3 g, 25.2 mmol), iron powder (6.0 g, 107.4 mmol), acetic acid (100 mL), and water (1 mL) were stirred at room temperature overnight. The mixture was diluted with Et_2O (100 mL) and water (100 mL). The aqueous phase was adjusted to pH 5 with a 4 N NaOH solution. The combined organic layers were washed with an aqueous saturated NaCl solution (100 mL), dried (MgSO_4), and concentrated under reduced pressure. The residue was purified by column chromatography with 50% ethyl acetate / 50% hexanes. It afforded the desired product as a white solid (4.8 g, 86%). TLC (50% EtOAc/50% hexane) R_f 0.28; $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ 6.04 (bs, 2H), 6.47-6.51 (m, 2H), 6.91 (d, $J=6.1$ Hz, 2H), 7.22 (t, $J=8.4$ Hz, 1H), 8.30 (d, $J=6.4$ Hz, 2H).

Using similar methods to those described for the preparation of Intermediates B1-B7, the following additional compounds were also prepared:

Table 2
Arylamine Intermediates B




Intermediate No.	Z	(R ₅) _n	-X-	A	Physical Properties
B8	CH	H	(4)-S-CH ₂ -	pyrid-4-yl	TLC R _f = 0.12 (50% EtOAc/50% hexanes). 1H NMR (DMSO- <i>d</i> ₆) δ 3.91 (s, 2H), 5.26 (bs, 2H), 6.44 (d, J = 8.7 Hz, 2H), 6.96 (d, J = 8.7 Hz, 2H), 7.12 (d, J = 6.3 Hz, 2H), 8.40 (d, J = 6.0 Hz, 2H).
B9	CH	3-CF ₃	(4)-S-	pyrid-4-yl	TLC R _f = 0.10 (50% EtOAc/50% hexanes). 1H NMR (DMSO- <i>d</i> ₆) δ 6.21 (bs, 2H), 6.84-6.87 (m, 3H), 7.10 (d, J = 2.4 Hz, 1H), 7.39 (d, J = 8.4 Hz, 1H), 8.29 (d, J = 6.3 Hz, 2H).
B10	CH	H	(4)-O-	isoquinolin-5-yl	
B11	CH	3-F	(4)-O-	isoquinolin-5-yl	1H NMR (DMSO- <i>d</i> ₆) δ 5.42 (bs, 2H), 6.41-6.55 (m, 2H), 6.81-7.05 (m, 2H), 7.48-7.54 (m, 1H), 7.73-7.76 (m, 1H), 8.06-8.08 (m, 1H), 8.54-8.56 (m, 1H), 9.32 (s, 1H).
B12	CH	3,5-(Cl) ₂	(4)-O-	isoquinolin-5-yl	TLC R _f = 0.29 (45% EtOAc/55% hexanes). 1H NMR (DMSO- <i>d</i> ₆) δ 5.73 (bs, 2H), 6.69 (dd, J = 1.1 and 8.0 Hz, 1H), 6.75 (s, 2H), 7.51 (t, J = 7.7 Hz, 1H), 7.78 (d, J = 8.2 Hz, 1H), 8.12 (d, J = 5.9 Hz, 1H), 8.58 (d, J = 5.6 Hz, 1H), 9.34 (bs, 1H).
B13	CH	H	(4)-S-	pyrid-4-yl	TLC R _f = 0.07 (100% EtOAc). 1H NMR (DMSO- <i>d</i> ₆) δ 5.84 (bs, 2H), 6.95-6.99 (m, 3H), 7.32 (d, J = 8.6 Hz, 1H), 8.00 (d, J = 2.8 Hz, 1H), 8.31 (d, J = 4.7 Hz, 2H).
B14	CH	3,5-(Cl) ₂	(4)-S-	pyrid-4-yl	1H NMR (DMSO- <i>d</i> ₆) δ 6.30 (bs, 2H), 6.82 (s, 2H), 6.89, (d, J = 6.0 Hz, 2H), 8.33 (d, J = 6.1 Hz, 2H).
B15	CH	2,5-(F) ₂	(4)-S-	pyrid-4-yl	
B16	CH	3-Cl	(4)-S-	pyrid-4-yl	
B17	CH	H	(4)-S-	isoquinolin-5-yl	
B18	CH	H	(4)-CH ₂ -S-	pyrid-4-yl	

Intermediate No.	Z	(R ₅) _n	-X-	A	Physical Properties
B19	CH	H	(4)-S-	pyrid-3-yl	
B20	CH	H	(3)-S-	pyrid-4-yl	
B21	CH	H	(4)-O-	quinolin-5-yl	

Intermediate C1


1-Chloro-2,3-difluoro-5-nitrobenzene



The compound can be obtained by oxidizing 3-chloro-4,5-difluoroaniline, described in JP 05059067, with hydrogen peroxide in trifluoroacetic acid according to a process described in Heaton, A. et al., J. Fluorine Chem. 1997, 81 (2), 133-138 and Krapcho, A.P. et al., J. Org. Chem. 1990, 55 (21), 5662-5664 for the preparation of analogous derivatives.

Intermediate C2

4-[(2-Fluoro-4-nitrophenyl)sulphanyl]pyridine



21 g (188.9 mmol) of 4-mercaptopyridine, 30.05 g (188.9 mmol) of 3,4-difluoro-nitrobenzene and 60.05 g (434.5 mmol) of potassium carbonate are dissolved in dimethylformamide, and the mixture is stirred at 40 °C for 3 hours. The reaction solution

is then diluted with 500 mL of ethyl acetate and 300 mL of water. The aqueous phase is extracted five times with in each case 100 mL of ethyl acetate. The combined organic phases are washed with 200 mL of saturated sodium chloride solution, dried over sodium sulphate and concentrated under reduced pressure using a rotary evaporator. The residue is purified by MPLC (mobile phase: ethyl acetate:cyclohexane 1:1).


This gives 37.3 g (79% of theory) of product.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6): δ = 7.28 (dd, 2H), 7.79 (t, 1H), 8.15 (dd, 1H), 8.30 (dd, 1H), 8.50 (dd, 2H)

LC-MS (method 4): RT = 2.68 min


MS (ESIpos): m/z = 251 ($\text{M}+\text{H}$)⁺

Intermediate C3 below can be prepared from the appropriate mercapto or hydroxy heterocycles and their corresponding 4-fluoro- and 4-chloronitrobenzene derivatives, analogously to the procedure described for C2.

C3		$^1\text{H-NMR}$ (300 MHz, DMSO-d_6): δ = 7.15 (dd, 2H), 8.37 (dd, 1H), 8.41-8.45 (m, 3H) HPLC (method 1): RT=3.77 min MS (CIpos): m/z =302 ($\text{M}+\text{NH}_4$) ⁺
----	---	--

Intermediate C4

3-Fluoro-4-(4-pyridinylsulphanyl)aniline




37 g (147.9 mmol) of C2 are dissolved in 1000 mL of ethanol, and 143.86 mL (2.95 mol) of hydrazine hydrate and 4 g of palladium on carbon are added. The reaction mixture is stirred under reflux overnight. After cooling, the mixture is filtered off with suction through silica gel, and the filter cake is washed with ethanol. The filtrate is concentrated under reduced pressure using a rotary evaporator. The residue is suspended in diethyl ether and filtered off with suction. The precipitate is then suspended in water and filtered off with suction. The product is washed two more times with a little water. Drying under high vacuum gives 27.3 g (84% of theory) of product.

$^1\text{H-NMR}$ (300 MHz, DMSO-d_6): δ = 6.02 (br.s, 2H), 6.49-6.54 (m, 2H), 6.93 (dd, 2H), 7.23 (t, 1H), 8.32 (dd, 2H)

LC-MS (method 4): RT 0.96 min


MS (ESIpos): m/z = 221 ($\text{M}+\text{H}^+$)

Intermediate C5 can be prepared analogously to the procedure described for C4 from the appropriate starting materials.

C5		<p>$^1\text{H-NMR}$ (200 MHz, DMSO-d_6): δ = 5.46 (br.s, 2H), 6.46 (d, 1H), 6.55 (dd, 1H), 6.85 (d, 1H), 7.04 (t, 1H), 7.54 (t, 1H), 7.77 (d, 1H), 8.10 (d, 1H), 8.58 (d, 1H), 9.35 (s, 1H)</p> <p>LC-MS (method 4): RT=1.95 min MS (ESIpos): m/z=255 ($\text{M}+\text{H}^+$)</p>
----	---	--

Intermediate C6

3-Chloro-5-fluoro-4-(4-pyridinylsulphanyl)aniline



3.19 g (11.205 mmol) of C3 are dissolved in 200 mL of ethanol. 638 mg (2.81 mmol) of platinum(IV) oxide are then added, and the mixture is stirred at RT and atmospheric pressure under an atmosphere of hydrogen for 2 hours. For work-up, the reaction solution is filtered off with suction through kieselguhr and washed thoroughly with ethanol. The filtrate is concentrated under reduced pressure.


This gives 2.755 g (81% of theory) of product.

¹H-NMR (200 MHz, DMSO-d₆): δ = 6.37 (s, 2H), 6.49 (dd, 1H), 6.72-6.74 (m, 1H), 6.93 (dd, 2H), 8.34 (dd, 2H)

HPLC (method 1): RT = 3.68 min

MS (ESIpos): m/z = 255 (M+H)⁺


Intermediate C7: 3,5-difluoro-4-(4-pyridinylsulfanyl)aniline



is synthesized analogously to C6 from 4-[(2,6-difluoro-4-nitrophenyl)sulfanyl]pyridine by catalytic reduction with hydrogen on Platin(IV)oxide.

¹H-NMR (300 MHz, DMSO-d₆): δ = 6.38 (br. s, 2H), 6.40 (m, 2H), 6.98 (dd, 2H), 8.35 (dd, 2H)


HPLC (method 1): RT = 3.56 min

4-[(2,6-difluoro-4-nitrophenyl)sulfanyl]pyridine

is synthesized analogously to C2 from 4-mercaptopyridine and 1,2,3-trifluoro-5-nitrobenzene.

$^1\text{H-NMR}$ (200 MHz, DMSO-d_6): δ = 7.22 (dd, 2H), 8.32 (mc, 2H), 8.44 (dd, 2H)


HPLC (method 1): RT = 3.64 min

1,2,3-trifluoro-5-nitrobenzene

is synthesized analogously to C1 from 3,4,5-trifluor-aniline by oxidation with hydrogenperoxide in trifluoroacetic acid.

GC-MS (method 13): RT = 3.15 min


MS (ESIpos): m/z = 177 (M^+)

Intermediate C8: 3-chloro-4-(4-pyridinylsulfanyl)aniline

is synthesized analogously to C6 from 4-[(2-chloro-4-nitrophenyl)sulfanyl]pyridine by catalytic reduction with hydrogen on Platin(IV)oxide.

LC-MS (method 7): RT = 2.70 min

MS (ESIpos): m/z = 237 (M+H)⁺

4-[(2-chloro-4-nitrophenyl)sulfanyl]pyridine

is synthesized analogously to C2 from 4-mercaptopyridine and 1,2-dichloro-4-nitrobenzene.

¹H-NMR (200 MHz, DMSO-d₆): δ = 7.42 (dd (2H), 7.55 (d, 1H), 8.19 (dd (1H), 8.47 (d, 1H), 8.50 (dd, 2H)

HPLC (method 1): RT = 3.72 min

HPLC, LCMS and GCMS methods:**Method 1 (HPLC):**

Instrument: HP 1100 with DAD detection; column: Kromasil RP-18, 60 mm × 2 mm, 3.5 µm; mobile phase: A=5 mL of HClO₄/l of H₂O, B=acetonitrile; gradient: 0 min 2% B, 0.5 min 2% B, 4.5 min 90% B, 6.5 min 90% B; flow rate: 0.75 mL/min; temp.: 30°C; detection UV 210 nm.

Method 7 (LCMS):

Instrument: Micromass Platform LCZ, HP1100; column: Symmetry C18, 50 mm × 2.1 mm, 3.5 µm; mobile phase A: water + 0.05% formic acid, mobile phase B: acetonitrile + 0.05% formic acid; gradient: 0.0 min 90% A → 4.0 min 10% A → 6.0 min 10% A; oven: 40°C; flow rate: 0.5 mL/min; UV detection: 208-400 nm.

Method 10 (LCMS):

Instrument: Waters Alliance 2790 LC; column: Symmetry C18, 50 mm × 2.1, 3.5 µm; mobile phase A: water + 0.1% formic acid, mobile phase B: acetonitrile + 0.1% formic acid; gradient: 0.0 min 5% B → 5.0 min 10% B → 6.0 min 10% B; temperature: 50°C; flow rate: 1.0 mL/min; UV detection: 210 nm.


Method 12 (LCMS):

Instrument: Micromass Platform LCZ, HP 1100; column: Grom-SIL120 ODS-4 HE, 50 mm × 2.0 mm, 3 µm; mobile phase A: 1l water + 1 mL 50% formic acid, mobile phase B: 1l acetonitrile + 1mL 50% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1 min 10% A → 4.5 min 10% A; temperature: 55°C; flow rate: 0.8 mL/min; UV detection: 208-400 nm


Method 13 (GCMS):

Instrument: Micromass GCT, GC6890; column: Restek RTX-35MS, 30 m × 250 µm × 0.25 µm; carrier gas: helium; flow rate: 0.88 mL/min; initial temperature: 60°C; front injector temp.: 250°C; gradient: 60°C (1 min const.), 16°C/min up to 300°C then 1 min const. 300°C.


The following additional compounds may be produced analogously to the relevant examples above with the (+) and (-) enantiomers being separated on a chiral column:




anti isomer, racemic mixture



anti isomer, enantiomer 1



anti isomer, enantiomer 2




anti isomer, racemic

(+) and (-) enantiomers of the above compound may be separated on a chiral column.

D. Preparation of Examples of the Invention

Example 1








Preparation of 4-({4-[(2-amino-6-ethyl-4-pyrimidinyl)amino]phenyl}sulfanyl)phenol



2-Amino-4-chloro-6-ethylpyrimidine (A8, 50 mg, 0.23 mmol) and Intermediate B2(40 mg, 0.25 mmol) were suspended in a mixture of 0.01 M aqueous HCl (230 μ L) and 1-butanol (230 μ L). The mixture was refluxed overnight. The reaction was cooled to room temperature and quenched with NaHCO₃ (satd eq). The precipitated solid was filtered and subsequently purified by reversed phase chromatography on a YMC Pack-pro C18 column (trademark) eluting with acetonitrile/H₂O (10:90 – 90:10 gradient). Desired compound (40.5 mg, 0.11 mmol, 51 % yield); mp = 181-184°C, TLC (95:5 CH₂Cl₂/MeOH); R_f = 0.13

Examples 2-14

Using the above procedures, the following examples of pyridines were synthesized and are summarized in Table 3.

Table 3

Ex. No.	Intermediate Pyrimidone (A)	Intermediate Amine (B)	R ₁	R ₂	R ₄	Analytical Data
2	A1	B1	(CH ₃) ₃ C-	H	a	¹ H NMR (DMSO- <i>d</i> ₆) δ 11.75 (s, 1H), 10.59 (s, 1H), 8.78 (d, <i>J</i> = 5.4, 2H), 8.21 (s, 1H), 7.87 – 7.97 (m, 5H), 7.47 (d, <i>J</i> = 8.1, 1H), 6.85 (d, <i>J</i> = 3.4, 1H), 6.21 (s, 1H), 1.29 (s, 9H); ES MS [M+H] ⁺ = 359.
3	A1	B1	-(CH ₂) ₅ -		a	mp = 227 – 230 °C; ES MS [M+H] ⁺ = 371; TLC: R _f = 0.21 (CH ₂ Cl ₂ -MeOH, 95:5).
4	A2	B1	-CH ₂ CH ₂ N(Bn)CH ₂ -		a	mp = 227 – 230 °C; ES MS (M+H) ⁺ = 448; TLC: R _f = 0.45 (CH ₂ Cl ₂ -MeOH, 95:5).
5	A1	B1	-(CH ₂) ₅ -		b	mp = 220 – 224 °C, ES MS [M+H] ⁺ = 364; TLC: R _f = 0.39 (CH ₂ Cl ₂ -MeOH, 95:5).
6	A1	B2	CF ₃	H	c	¹ H NMR (Methanol- <i>d</i> ₄) δ 8.48 – 8.50 (m, 2H), 8.33 (dd, <i>J</i> = 1.1, 1.4, 1H), 7.60 – 7.62 (m, 4H), 6.45 (s, 1H); ES MS [M+H] ⁺ = 382; TLC: R _f = 0.57 (CH ₂ Cl ₂ -MeOH, 90:10).


Ex. No.	Intermediate Pyrimidone (A)	Intermediate Amine (B)	R ₁	R ₂	R ₄	Analytical Data
7	A1	B1	-(CH ₂) ₃ -		c	¹ H NMR (Methanol-d ₄) δ 8.30 (d, J = 7.1, 2H), 7.71 (t, J = 7.8, 1H), 7.29 – 7.38 (m, 4H), 3.02 (t, J = 7.7, 2H), 2.87 (t, J = 5.5, 2H), 2.28 (quint, J = 7.6, 2H); ES MS [M+H] ⁺ = 356; TLC: R _f = 0.34 (CH ₂ Cl ₂ -MeOH, 90:10).
8	Aldrich	B1	H	H	d	¹ H NMR (Acetone-d ₆) δ 7.80 (d, J = 7.1, 1H), 7.70 (s, 2H), 7.34 (d, J = 8.4, 2H), 7.10 (d, J = 8.8, 2H), 6.91 (d, J = 8.2, 2H), 6.36 (d, J = 7.1, 1H) TLC: R _f = 0.07 (95:5 CH ₂ Cl ₂ /MeOH).
9	A1	B1	CH ₃ CH ₂ -	H	d	mp = 181-184°C, TLC: R _f = 0.13 (95:5 CH ₂ Cl ₂ /MeOH).
10	A1	B1	(CH ₃) ₂ CH-	H	d	mp = 86-90°C, TLC: R _f = 0.10 (95:5 CH ₂ Cl ₂ /MeOH).
11	A1	B1	(CH ₃) ₃ C-	H	d	mp = 95-99°C, TLC: R _f = 0.16 (95:5 CH ₂ Cl ₂ /MeOH).

Ex. No.	Intermediate Pyrimidone (A)	Intermediate Amine (B)	R ₁	R ₂	R ₄	Analytical Data
12	A1	B1	CH ₃ -	Cl-	d	mp = 242-245°C, TLC: R _f = 0.32 (95:5 CH ₂ Cl ₂ /MeOH).
13	A1	B1	CH ₃ -	H	e	mp = 209-212°C, TLC: R _f = 0.08 (95:5 CH ₂ Cl ₂ /MeOH).
14	A1	B1	CH ₃ -	H	f	mp = 271-274°C, TLC: R _f = 0.05 (95:5 CH ₂ Cl ₂ /MeOH).

By selecting combinations of the appropriate Intermediates A1-A21 with Intermediates B1-B17, a variety of products were prepared in like manner and are described in Examples 15-18

Example 15


Preparation of *N*-(2-amino-6-methyl-4-pyrimidinyl)-*N*-[4-(3-pyridinylsulfanyl)phenyl]amine



Prepared in 5% yield from A7 and B19: TLC (7% MeOH in CH₂Cl₂) R_f 0.49; MS (ES) 310 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 2.13 (s, 3H), 5.94 (s, 1H), 6.48 (bs, 2H), 7.31-7.39 (m, 3H), 7.56 (td, *J*=1.6, 8.2 Hz, 1H), 7.81 (d, *J*=8.9 Hz, 2H), 8.38-8.40 (m, 2H), 9.50 (bs, 1H).

Example 16


Preparation of *N*-(2-amino-6-methyl-4-pyrimidinyl)-*N*-[3-(4-pyridinylsulfanyl)phenyl]amine



Prepared in 29% yield from A7 and B20: TLC (6% MeOH in CH₂Cl₂) R_f 0.37; MS (ES) 310 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 2.08 (s, 3H), 5.86 (s, 1H), 6.17 (bs, 2H), 7.06 (d, *J*=6.0 Hz, 2H), 7.09 (d, *J*=7.5 Hz, 1H), 7.39 (t, *J*=7.8 Hz, 1H), 7.78 (t, *J*=2.2 Hz, 1H), 7.99 (dd, *J*=1.4, 8.1 Hz, 1H), 8.36 (d, *J*=6.3 Hz, 2H), 9.18 (bs, 1H).

Example 17


Preparation of *N*-(2-anilino-6-methyl-4-pyrimidinyl)-*N*-[4-(4-pyridinylsulfanyl)phenyl]amine



Prepared in 1% yield from A16 and B5: TLC (50% EtOAc/50% hexanes) R_f 0.15; MS (ES) 386 $[M+H]^+$; $^1\text{H-NMR}$ (DMSO- d_6) δ 2.25 (s, 3H), 6.14 (s, 1H), 6.89 (t, $J=7.4$ Hz, 1H), 6.97 (d, $J=5.7$ Hz, 2H), 7.23 (t, $J=7.2$ Hz, 2H), 7.47 (d, $J=9.0$ Hz, 2H), 7.71 (dd, $J=0.9$, 8.6 Hz, 2H), 7.89 (d, $J=8.5$ Hz, 2H), 8.33 (bs, 2H), 9.21 (s, 1H), 9.55 (s, 1H).

Example 18

Preparation of *N*-(2-amino-6-methyl-4-pyrimidinyl)-*N*-[4-(5-quinolinylthio)phenyl]amine



Prepared in 71% yield from A7 and B21: TLC (6% MeOH in CH_2Cl_2) R_f 0.34; MS (ES) 344 $[M+H]^+$; $^1\text{H-NMR}$ (DMSO- d_6) δ 2.08 (s, 3H), 5.85 (s, 1H), 6.11 (bs, 2H), 6.88 (d, $J=7.3$ Hz, 1H), 7.04 (d, $J=8.6$ Hz, 2H), 7.56 (dd, $J=4.0$, 8.5 Hz, 1H), 7.65 (t, $J=8.5$ Hz, 1H), 7.73-7.76 (m, 3H), 8.57 (d, $J=8.6$ Hz, 1H), 8.94 (d, $J=3.9$ Hz, 1H), 9.03 (bs, 1H).


Examples 19-22

Using the following general method, Examples 19-22 were prepared

A suspension of Intermediate A (1.0 mmol), Intermediate B (1.0 mmol), and HCl (0.1 mL) in H_2O (1.0 mL) was heated to 70 °C in a 5 mL reaction vial overnight. The reaction mixture was diluted with MeOH, treated with saturated NaHCO_3 , coated on silica and purified by MPLC (Biotage) with 5% MeOH in CH_2Cl_2 .

Example 19


Preparation of *N*-(2-amino-6-phenyl-4-pyrimidinyl)-*N*-[3,5-dichloro-4-(4-pyridinylsulfanyl)phenyl]amine



Prepared in 40% yield from A17 and B14: TLC (4% MeOH in CH₂Cl₂) R_f 0.42; MS (ES) 440 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 6.53 (s, 1H), 6.67 (bs, 2H), 6.94 (d, *J*=6.1 Hz, 2H), 7.47-7.51 (m, 3H), 7.93-7.96 (m, 2H), 8.18 (s, 2H), 8.36 (d, *J*=5.7 Hz, 2H), 9.88 (bs, 1H).

Example 20


Preparation of *N*-[2-amino-6-(3-nitrophenyl)-4-pyrimidinyl]-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine



Prepared in 97% yield from A20 and B7: TLC (50% EtOAc/50% hexanes) R_f 0.15; MS (ES) 435 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 6.69 (s, 1H), 6.78 (bs, 2H), 6.98 (d, *J*=4.8 Hz, 2H), 7.46-7.57 (m, 2H), 7.79 (t, *J*=7.9 Hz, 1H), 8.33-8.35 (m, 5H), 8.79 (s, 1H), 9.94 (bs, 1H).

Example 21


Preparation of *N*-[2-amino-6-(2-furyl)-4-pyrimidinyl]-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine



Prepared in 12% yield from A4 and B7: TLC (6% MeOH in CH₂Cl₂) R_f 0.37; MS (ES) 380 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 6.45 (s, 1H), 6.58 (bs, 2H), 6.64 (dd, *J*=1.9, 3.5 Hz, 1H), 6.97-6.70 (m, 3H), 7.44 (dd, *J*=1.9, 8.4 Hz, 1H), 7.53 (t, *J*=8.6 Hz, 1H), 7.84 (d, *J*=1.6 Hz, 1H), 8.29-8.35 (m, 3H), 9.80 (bs, 1H).

Example 22

Preparation of N-[2-amino-6-(2-thienyl)-4-pyrimidinyl]-N-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine




Prepared in 29% yield from A3 and B7. TLC (4% MeOH in CH₂Cl₂) R_f 0.22; MS (ES) 396 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 6.47 (s, 1H), 6.61 (bs, 2H), 6.98 (d, *J*=6.4 Hz, 2H), 7.15-7.18 (m, 1H), 7.43-7.68 (m, 4H), 8.29-8.36 (m, 3H), 9.79 (bs, 1H).

Using analogous methods, Example 23 is synthesized from 4-chloro-6-furan-3-yl-2-methyl-pyrimidine and Intermediate B7.

Example 23

Preparation of 6-furan-3-yl-N⁴-[4-(pyridin-4-yl-sulfanyl)-phenyl]-pyrimidine-2,4-diamine




Examples 24-26

The following general procedure and the appropriate Intermediates A and B provided the compounds described in Table 4 below.

A mixture of Intermediate B (0.250 mmol) and Intermediate (0.25 mmol) in 0.01 M aqueous HCl (500 μ L) was refluxed for 6 h. The reaction was cooled to room temperature and the solvent was evaporated by vacuum. The residue was purified by reverse phase chromatography on a YMC Pack-pro[®] C18 column eluting with acetonitrile/H₂O (10:90 – 90:10 gradient) to give the desired compound.

In some cases, the compound was further purified by preparative TLC eluting with CH₂Cl₂-MeOH (90:10) to give the desired product.

Table 4



Ex. No.	Yield %	A int.	B int.	R ₅	R ₁	R ₂	Note
24	10	A13	B5	H	-(CH ₂) ₅ -		1
25	8	A6	B7	F	CF ₃	H	2
26	33	A14	B7	F	-(CH ₂) ₃ -		3


1) mp = 220 – 224 °C, ES MS [M+H]⁺ = 364; TLC (CH₂Cl₂-MeOH, 95:5; R_f = 0.39.

2) ¹H NMR (Methanol-*d*₄) δ 8.48 – 8.50 (m, 2H), 8.33 (dd, *J* = 1.1, 1.4, 1H), 7.60 – 7.62 (m, 4H), 6.45 (s, 1H); ES MS [M+H]⁺ = 382; TLC (CH₂Cl₂-MeOH, 90:10); R_f = 0.57.


3) ¹H NMR (Methanol-*d*₄) δ 8.30 (d, *J* = 7.1, 2H), 7.71 (t, *J* = 7.8, 1H), 7.29 – 7.38 (m, 4H), 3.02 (t, *J* = 7.7, 2H), 2.87 (t, *J* = 5.5, 2H), 2.28 (quint, *J* = 7.6, 2H); ES MS [M+H]⁺ = 356; TLC: R_f = 0.34 (CH₂Cl₂-MeOH, 90:10).

Example 27

Preparation of *N*-[2-amino-6-(4-ethoxyphenyl)-4-pyrimidinyl]-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine




Step 1. Preparation of *N*-(2-amino-6-chloro-4-pyrimidinyl)-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine



2-Amino-4,6 dichloropyrimidine (A21, 12 mmol) and 3-fluoro-4-(4-pyridinylthio)-aniline (B7, 12 mmol) were suspended in water (150 mL) and treated with 10 drops of concentrated hydrochloric acid. The mixture was stirred at 100 °C overnight. The clear solution was then neutralized with ammonium hydroxide. The precipitated yellow product was filtered, washed with water, and purified by column chromatography with 1-3% MeOH in CH₂Cl₂ to give the desired product as a white solid (1.98 g, 47%).

Step 2. Preparation of *N*-[2-amino-6-(4-ethoxyphenyl)-4-pyrimidinyl]-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine



A solution of *N*-(2-amino-6-chloro-4-pyrimidinyl)-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine from step 1 in toluene (2 mL) in ethanol (1 mL) was treated with aqueous sodium carbonate solution (2M, 0.5 mL). The mixture was then treated with 4-ethoxyphenylboronic acid (0.36 mmol), tetrakis(triphenylphosphine)-palladium (0.009 mmol), and allowed to stir at 120 °C under argon overnight. At this time, the solvents were removed at reduced pressure and the residue was purified by preparative silica gel with 3% MeOH in CH₂Cl₂ to afford the desired product as a white solid (21 mg).


Parallel synthesis methods

Example 28-33

An equivalent of *N*-(2-amino-6-chloro-4-pyrimidinyl)-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine (Example 26, step 1, 0.1 mmol) and four equivalents of the desired amine (R₆R₇NH) were suspended in *n*-butanol (1 mL) and shaken at 110 °C overnight to 2 days. The solvent was removed at reduced pressure and the residue was

purified either by preparative silica gel plate or by HPLC. The products were checked by LC-MS and TLC and are listed in Table 5 below.


Table 5



Ex No.	R ₆	R ₇
28	-(CH ₂ CH(Me)OCH(Me)CH ₂ -	
29		
30	<i>n</i> -Pr	<i>n</i> -Pr
31	-(CH ₂ CH ₂ OCH ₂ CH ₂ -	
32	<i>i</i> -Pr	H
33	cyc-Pent	H

Example 34

Preparation of *N*-[2-amino-6-(3-aminophenyl)-4-pyrimidinyl]-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine



A dry flask under argon was charged with 10% palladium on carbon (20 mg). A solution of *N*-[2-amino-6-(3-nitrophenyl)-4-pyrimidinyl]-*N*-[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine (Example 20, 200 mg, 0.46 mmol) in ethanol (12 mL) and EtOAc (1 mL) was added via syringe through a septa to the palladium on carbon. The flask was then fitted with a hydrogen balloon and stirred at room temperature for 48 h. At this time the palladium was filtered through a pad of Celite[®]. The filtrate was coated on silica and purified on the MPLC (Biotage) with 2-8% MeOH in CH₂Cl₂ to afford the desired product as a solid (120 mg, 64%). TLC (6% MeOH in CH₂Cl₂) R_f 0.26; MS (ES) 405 [M+H]⁺; ¹H-NMR (DMSO-*d*₆) δ 5.22 (bs, 2H), 6.45 (s, 1H), 6.49 (bs, 2H), 6.62-6.65 (m, 1H), 6.98 (d, *J*=6.4 Hz, 2H), 7.02-7.16 (m, 3H), 7.43-7.55 (m, 2H), 8.30-8.35 (m, 3H), 9.75 (bs, 1H).

Examples 35 to 41 listed in the table below can be prepared analogously to the procedure described in Example 1 using the following starting materials:




Example	starting material 1	starting material 2
35	B7 or C4	2,4-diamino-6-chloro-pyrimidine
36	B7 or C4	2-amino-4-chloro-6- <i>N</i> -methylamino-pyrimidine *
37	C5	2,4-diamino-6-chloro-pyrimidine
38	C7	A 21
39	C8	A 21
40	C6	A 2
41	C7	A 2





* was internally available, cited in:

DE 839640;

US 2,585,906;

Autenrieth et al., Justus Liebigs Ann. Chem., 1977, 1194,1210.

Example	Structure	Analytical data
35		¹ H-NMR (300 MHz, DMSO-d ₆) δ = 5.22 (s, 1H), 5.82 (d, 2H), 5.95 (s, 2H), 6.97 (d, 2H), 7.35-7.46 (m, 2H), 8.16 (dd, 1H), 8.34 (d, 2H), 9.12 (s, 1H) LC-MS (method 7): RT = 0.43 min MS (ESIpos): m/z = 329 (M+H) ⁺
36		¹ H-NMR (200 MHz, DMSO-d ₆) δ = 2.78 (d, 3H), 5.19 (s, 1H), 5.76 (s, 1H), 5.92 (s, 2H), 6.96 (d, 2H), 7.32-7.48 (m, 2H), 8.20 (dd, 1H), 8.75 (d, 2H), 9.19 (s, 1H). MS (ESIpos): m/z = 343
37		LC/MS (method 10): RT = 1.37 min. MS (ESI pos.): m/z = 363 (M+H) ⁺ ,


Example	Structure	Analytical data
38		MS (ESI pos.): $m/z = 366 (M+H)^+$, HPLC (method 1): RT = 3.75 min.
39		LC/MS (method 12): RT = 2.90 min. MS (ESI pos.): $m/z = 363 (M+H)^+$,
40		MS (ESI pos.): $m/z = 425 (M+H)^+$, HPLC (method 1): RT = 3.45 min.
41		LC/MS (method 10): RT = 1.59 min. MS (ESI pos.): $m/z = 409 (M+H)^+$, $205 (M+H)^{2+}$

Rho Kinase Biochemical Assay:

ROCK-1 activity criteria: 0 no effect (< 40% inhibition), 1 effect (> 40% inhibition). The assay tests for inhibition of ROCK-1 phosphorylation of MBP (Myelin Basic Protein). The reaction (100 µl final volume) is carried out in polypropylene 96-well plates in 50 mM HEPES buffer pH 7.5 containing 5 mM MgCl₂ and 1 mM DTT. For each well, gstROCK1 (0.25 µgs of BAYER DRT gstROCK1) is combined with MBP (1 µg) in reaction buffer (70 µL combined volume). Inhibitors (5 µL of 20x conc. in 40% DMSO) are added to each well to give an 8 point dose response range from 1.0 µM to .5 nM. The reaction is begun by adding 25 µL of ATP (4x = 12 µM) in reaction buffer containing 0.8 µCi of ³³P gamma-ATP (4x) for a final concentration of 3 µM cold and 0.2 µCi hot ATP. Plates were incubated for 1 hour at room temperature with the reaction being stopped by addition of 7 µL of 1 N HCl. The radioactively labeled MBP was transferred to P30 filtermats (EG&G Wallac), washed in 1% phosphoric acid followed by brief washes in water. The filtermats were then dried and the incorporation of ³³P detected by liquid scintillation counting. Background ³³P incorporation is determined by ROCK1 autophosphorylation without MBP. The data are expressed as percent inhibition: % inhibition = 1-((cpm with inhibitor-background)/(cpm without inhibitor - background)) * 100.

WHAT WE CLAIM:

1. A compound of formula I




wherein R_1 and R_2 are each independently

H, halo, alkyl optionally substituted by halo up to perhalo, cycloalkyl, alkenyl, alkynyl, NO_2 , NH_2 , NR_6R_7 , or furyl, thienyl, pyridyl, trifluoromethyl or phenyl each optionally substituted with NH_2 , NO_2 , trifluoromethyl or alkoxy; or

R_1 and R_2 may be taken together to form a ring of from 5 to 7 members optionally interrupted by N and optionally substituted on N by benzyl;

R_3 is NH_2 or $-\text{NH}-$ phenyl optionally substituted with halo, C_1-C_4 alkyl, trifluoromethyl, nitro or amino;

R_4 is



X is a linker substituted at the 3 or 4 position of the ring and is

O, S, $-\text{S}-\text{CH}_2-$, $-(\text{CH}_2)_m-$, or $-(\text{C}=\text{O})-$;

A is phenyl optionally substituted with alkylthio or OH,

pyridyl,

quinolyl or

isoquinolyl;

each R_5 independently is halo, alkyl optionally substituted by halo up to perhalo, cycloalkyl, alkoxy, alkenyl, alkynyl, NO_2 , NH_2 , or trifluoromethyl;

n is 0,1,2,3 or 4;




m is 1 or 2; and





R₆ and R₇ are each independently H, alkyl, cycloalkyl, or phenyl optionally substituted with halo, CF₃, alkyl, nitro or amino; or




R₆ and R₇ may form, together with the N atom to which they are attached, a heterocyclic ring optionally substituted with alkyl, optionally interrupted by O, or optionally fused to phenyl;





or a pharmaceutically acceptable salt thereof,





with the proviso that formula I does not include the following compounds:
























































or




2. A compound of formula I according to claim 1, wherein

R_1 and R_2 are each independently H, halo, C_{1-12} -alkyl optionally substituted by halo up to perhalo, C_{2-12} , alkenyl, C_{2-12} -alkynyl, NO_2 , NH_2 , NR_6R_7 , alkyl, furyl, thienyl, pyridyl, phenyl optionally substituted with NO_2 , trifluoromethyl or NR_6R_7 ; or

R_1 and R_2 together form a ring of from 5 to 7 members optionally interrupted by N and optionally substituted on N by benzyl;

R_3 is NH_2 or $-NH-$ phenyl optionally substituted with halo, C_1-C_4 alkyl, trifluoromethyl, nitro or amino;

R_4 is



X is a linker substituted at the 3 or 4 position of the ring and is selected from

O, S, $-\text{S}-\text{CH}_2-$, $-\text{CH}_2-\text{S}-$, $-(\text{CH}_2)_m-$, or $-(\text{C}=\text{O})-$;

A is phenyl optionally substituted by C_{1-4} alkylthio or OH,

pyridyl,

quinolyl or

isoquinolyl;

each R_5 independently is halo, C_{1-12} alkyl, optionally substituted by halo, up to the perhalo level, C_{2-12} -alkenyl, C_{2-12} -alkynyl, NO_2 , NH_2 or trifluoromethyl;

n is 0, 1, 2, 3 or 4;

m is 1 or 2; and

R_6 and R_7 are each independently H, C_1-C_6 alkyl, or phenyl optionally substituted with halo, CF_3 , C_1-C_4 alkyl, nitro or amino.

3. A compound according to claim 1, wherein X is not S or A is not pyridyl.

4. A compound according to claim 1, wherein X is not S or $-\text{S}-\text{CH}_2-$.





5. A compound according to claim 1, wherein n is 1-4.




6. A compound according to claim 1, wherein n is 1-4 and each R_5 is independently halo, C_3-C_{12} alkyl optionally substituted by halo up to the perhalo level, C_{2-12} -alkenyl, C_{2-12} -alkynyl, NO_2 , NH_2 or trifluoromethyl.

7. A compound according to claim 1, which is 6-*tert*-butyl- N^4 -(1-pyridin-4-yl-1*H*-indol-5-yl)-pyrimidine-2,4-diamine, N^4 -(1-pyridin-4-yl-1*H*-indol-5-yl)-6,7,8,9-tetrahydro-5*H*-cycloheptapyrimidine-2,4-diamine, 7-benzyl- N^4 -(1-pyridin-4-yl-1*H*-indol-5-yl)-5,6,7,8-tetrahydro-pyrido[3,4-*d*]pyrimidine-2,4-diamine, N^4 -[4-(pyridin-4-ylsulfanyl)-phenyl]-6,7,8,9-tetrahydro-5*H*-cycloheptapyrimidine-2,4-diamine, N^4 -[3-


fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-6-trifluoromethyl-pyrimidine-2,4-diamine, N^4 -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-6,7-dihydro-5*H*-cyclopentapyrimidine-2,4-diamine, 4-[4-(2-amino-pyrimidin-4-ylamino)-phenylsulfanyl]-phenol, 4-[4-(2-amino-6-ethyl-pyrimidin-4-ylamino)-phenylsulfanyl]-phenol, 4-[4-(2-amino-6-isopropyl-pyrimidin-4-ylamino)-phenylsulfanyl]-phenol, 4-[4-(2-amino-6-*tert*-butyl-pyrimidin-4-ylamino)-phenylsulfanyl]-phenol, 4-[4-(2-amino-5-chloro-6-methyl-pyrimidin-4-ylamino)-phenylsulfanyl]-phenol, [3-(2-amino-5-chloro-6-methyl-pyrimidin-4-ylamino)-phenyl]-(4-methylsulfanyl-phenyl)-methanone, 4-[4-(2-amino-5-chloro-6-methyl-pyrimidin-4-ylamino)-phenoxy]-phenol, N^4 -[4-(pyridin-4-ylsulfanyl)-phenyl]-6,7,8,9-tetrahydro-5*H*-cycloheptapyrimidine-2,4-diamine, N^4 -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-6-trifluoromethyl-pyrimidine-2,4-diamine, N^4 -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-6,7-dihydro-5*H*-cyclopentapyrimidine-2,4-diamine, 6-(2,6-dimethyl-morpholin-4-yl)- N^4 -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-pyrimidine-2,4-diamine, 6-(2,3-dihydro-indol-1-yl)- N^4 -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-pyrimidine-2,4-diamine, N -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]- N'' , N'' -dipropyl-pyrimidine-2,4,6-triamine, N^4 -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-6-morpholin-4-yl-pyrimidine-2,4-diamine, N -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]- N'' -isopropyl-pyrimidine-2,4,6-triamine, N -cyclopentyl- N'' -[3-fluoro-4-(pyridin-4-ylsulfanyl)-phenyl]-pyrimidine-2,4,6-triamine, 4-(4-[(2-amino-6-ethyl-4-pyrimidinyl)amino]phenyl)sulfanylphenol, N -(2-amino-6-methyl-4-pyrimidinyl)- N -[4-(3-pyridinylsulfanyl)phenyl]amine, N -(2-amino-6-methyl-4-pyrimidinyl)- N -[3-(4-pyridinylsulfanyl)phenyl]amine, N -(2-anilino-6-methyl-4-pyrimidinyl)- N -[4-(4-pyridinylsulfanyl)phenyl]amine, N -(2-amino-6-methyl-4-pyrimidinyl)- N -[4-(5-quinolinylloxy)phenyl]amine, N -(2-amino-6-phenyl-4-pyrimidinyl)- N -[3,5-dichloro-4-(4-pyridinylsulfanyl)phenyl]amine, N -[2-amino-6-(3-nitrophenyl)-4-pyrimidinyl]- N -[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine, N -[2-amino-6-(2-furyl)-4-pyrimidinyl]- N -[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine, N -[2-amino-6-(2-thienyl)-4-pyrimidinyl]- N -[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine, or N -[2-amino-6-(4-ethoxyphenyl)-4-pyrimidinyl]- N -[3-fluoro-4-(4-pyridinylsulfanyl)phenyl]amine.



8. A compound according to claim 1, of the formula








9. A compound according to claim 1, of the formula





10. A compound according to claim 1, of the formula





11. A method of treating an indication mediated by Rho-kinase, comprising administering a compound of claim 1.


12. A method of treating hypertension, atherosclerosis, restenosis, cerebral ischemia, cerebral vasospasm, neuronal degeneration, spinal cord injury, cancer of the breast, colon, prostate, ovaries, brain or lung, thrombotic disorders, asthma, glaucoma, osteoporosis or erectile dysfunction, comprising administering to host in need thereof a compound according to claim 1.

13. A process according to claim 11, wherein the host is a human.


14. A process according to claim 12, wherein the host is a human.

15. A process for the preparation of a compound of claim 1, comprising reacting

a)



or b)



and optionally, where R_1 is NR_6R_7 , reacting the chloropyrimidine of Formula Ia with an amine of Formula R_6R_7NH .

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D239/48 C07D413/12 C07D403/12 C07D413/14
 C07D409/14 C07D471/04 A61K31/506

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 01 53331 A (ALI ANMAR ;NI FENG (CA); CHEN ZHIGANG (CA); GOUR BARBARA J (CA); A) 26 July 2001 (2001-07-26) figure 23E; example 425</p> <p style="text-align: center;">--- -/--</p>	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

30 April 2003

Date of mailing of the international search report

19/05/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Stix-Malaun, E

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; BLACKBURN, THOMAS P. ET AL: "Preparation of pyrimidine and indol-2-one derivatives as galanin GAL3 receptor antagonists for the treatment of depression and/or anxiety" retrieved from STN Database accession no. 137:154941 XP002239813 abstract & WO 2002 060392 A (SYNAPTIC PHARMACEUTICAL CORPORATION, USA) 8 August 2002 (2002-08-08)</p>	1
A	<p>WO 97 19065 A (CELLTECH THERAPEUTICS LTD ;DAVIS PETER DAVID (GB); MOFFAT DAVID FE) 29 May 1997 (1997-05-29) abstract claims examples</p>	1-15
A	<p>WO 01 19802 A (TANABE SEIYAKU CO ;KIKKAWA KOHEI (JP); MATSUKI KENJI (JP); OMORI K) 22 March 2001 (2001-03-22) abstract claims examples</p>	1-15
A	<p>DE 198 51 421 A (BOEHRINGER INGELHEIM PHARMA) 11 May 2000 (2000-05-11) abstract claims examples</p>	1-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 03/01839

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 11-14 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Internati pplication No

PCT/US 03/01839

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0153331	A	26-07-2001	AU 3115401 A	31-07-2001
			WO 0153331 A2	26-07-2001
			US 2002168761 A1	14-11-2002
WO 2002060392	A		NONE	
WO 9719065	A	29-05-1997	AU 7631496 A	11-06-1997
			EP 0862560 A1	09-09-1998
			WO 9719065 A1	29-05-1997
			US 6235746 B1	22-05-2001
			US 5958935 A	28-09-1999
WO 0119802	A	22-03-2001	AU 7311800 A	17-04-2001
			BG 106566 A	28-02-2003
			BR 0014526 A	18-06-2002
			CA 2383466 A1	22-03-2001
			CN 1374953 T	16-10-2002
			EP 1219609 A1	03-07-2002
			WO 0119802 A1	22-03-2001
			JP 2002012587 A	15-01-2002
			NO 20021308 A	24-04-2002
			TR 200200701 T2	21-06-2002
			US 2003032647 A1	13-02-2003
			AU 4114201 A	12-11-2001
			EP 1277741 A1	22-01-2003
			WO 0183460 A1	08-11-2001
DE 19851421	A	11-05-2000	DE 19851421 A1	11-05-2000
			AU 1378500 A	29-05-2000
			WO 0027826 A1	18-05-2000